

A Rule Format for Unit Elements^{*}

Luca Aceto¹, Anna Ingólfssdóttir¹,
MohammadReza Mousavi², and Michel A. Reniers²

¹ ICE-TCS, School of Computer Science, Reykjavik University,
Kringlan 1, IS-103 Reykjavik, Iceland

² Department of Computer Science, Eindhoven University of Technology,
P.O. Box 513, NL-5600 MB Eindhoven, The Netherlands

Abstract. This paper offers a meta-theorem for languages with a Structural Operational Semantics (SOS) in the style of Plotkin. Namely, it proposes a generic rule format for SOS guaranteeing that certain constants act as left- or right-unit elements for a set of binary operators. We show the generality of our format by applying it to a wide range of operators from the literature on process calculi.

1 Introduction

In many process algebras and specification languages, one encounters constructs that are unit elements for certain composition operators. The concept of (left) unit element for a binary operator f can be concisely summarized in the following algebraic equation, where $\mathbf{0}$ is the left-unit element for f : $f(\mathbf{0}, x) = x$.

In this paper, we propose a generic rule format guaranteeing that certain constants are left- or right-unit elements for a set of binary operators, whose semantics is defined using Plotkin’s style of Structural Operational Semantics (SOS) [2, 12, 13]. The notions of left and right unit are defined with respect to a notion of behavioural equivalence. There are various notions of behavioural equivalence presented in the literature (see, e.g., [7]), which are, by and large, weaker than bisimilarity. Thus, to be as general as possible, we prove our main result for all equivalences that contain, i.e., are weaker than, bisimilarity.

This paper is part of our ongoing line of research on capturing basic properties of composition operators in terms of syntactic rule formats, exemplified by rule formats for commutativity [11], associativity [6], determinism and idempotence [1].

This line of research serves multiple purposes. Firstly, it paves the way for a tool-set that can mechanically prove such properties without involving user interaction. Secondly, it provides us with an insight as to the semantic nature of

^{*} The work of Aceto and Ingólfssdóttir has been partially supported by the projects “The Equational Logic of Parallel Processes” (nr. 060013021), and “New Developments in Operational Semantics” (nr. 080039021) of the Icelandic Research Fund. The first author dedicates the paper to the memory of his mother, Imelde Diomedea Aceto, who passed away a year ago.

such properties and its link to the syntax of SOS deduction rules. In other words, our rule formats may serve as a guideline for language designers who want to ensure, a priori, that the constructs under design enjoy certain basic algebraic properties.

The rest of this paper is organized as follows. In Section 2, we define some basic notions that are required for the technical developments in the rest of the paper. In Section 3, we present our rule format and prove that it guarantees the unit element property. In Section 4, we apply the rule format to various examples from the literature. In order to ease the application of our rule format to operators whose operational semantics is specified using predicates, we extend the format to that setting in Section 4.2. Section 5 concludes the paper and discusses directions for future work. Due to space limitations proofs are omitted.³

2 Preliminaries

We begin by recalling the basic notions from the theory of SOS that are needed in the remainder of this study. We refer the interested readers to, e.g., [2, 12] for more information and background.

Definition 1 (Signatures, Terms and Substitutions) *We let V represent an infinite set of variables and use $x, x', x_i, y, y', y_i, \dots$ to range over elements of V . A signature Σ is a set of function symbols, each with a fixed arity. We call these symbols operators and usually represent them by f, g, \dots . An operator with arity zero is called a constant. We define the set $\mathbb{T}(\Sigma)$ of terms over Σ as the smallest set satisfying the following constraints.*

- A variable $x \in V$ is a term.
- If $f \in \Sigma$ has arity n and t_1, \dots, t_n are terms, then $f(t_1, \dots, t_n)$ is a term.

We use s, t , possibly subscripted and/or superscripted, to range over terms. We write $t_1 \equiv t_2$ if t_1 and t_2 are syntactically equal. The function $\text{vars} : \mathbb{T}(\Sigma) \rightarrow 2^V$ gives the set of variables appearing in a term. The set $\mathbb{C}(\Sigma) \subseteq \mathbb{T}(\Sigma)$ is the set of closed terms, i.e., terms that contain no variables. We use p, q, p', p_i, \dots to range over closed terms. A substitution σ is a function of type $V \rightarrow \mathbb{T}(\Sigma)$. We extend the domain of substitutions to terms homomorphically and write $\sigma(t)$ for the result of applying the substitution σ to the term t . If the range of a substitution lies in $\mathbb{C}(\Sigma)$, we say that it is a closed substitution. An explicit substitution $[x \mapsto t]$ maps x to t and is the identity function on all variables but x .

Definition 2 (Transition System Specifications) *A transition system specification (TSS) is a triple (Σ, \mathcal{L}, D) where*

- Σ is a signature.

³ For reviewers' convenience, the omitted proofs are included in the appendices. Upon acceptance of the paper, a technical report with the proofs will appear and a reference to the technical report will be given in the paper.

- \mathcal{L} is a set of labels (or actions) ranged over by a, b, l . If $l \in \mathcal{L}$, and $t, t' \in \mathbb{T}(\Sigma)$ we say that $t \xrightarrow{l} t'$ is a positive transition formula and $t \xrightarrow{l} \neg$ is a negative transition formula. A transition formula (or just formula), typically denoted by ϕ or ψ , is either a negative transition formula or a positive one.
- D is a set of deduction rules, i.e., tuples of the form (Φ, ϕ) where Φ is a set of formulae and ϕ is a positive formula. We call the formulae contained in Φ the premises of the rule and ϕ the conclusion.

We write $\text{vars}(r)$ to denote the set of variables appearing in a deduction rule r . We say that a formula is closed if all of its terms are closed. Substitutions are also extended to formulae and sets of formulae in the natural way. For a rule r and a substitution σ , the rule $\sigma(r)$ is called a substitution instance of r . A set of positive closed formulae is called a transition relation.

We often refer to a positive transition formula $t \xrightarrow{l} t'$ as a *transition* with t being its *source*, l its *label*, and t' its *target*. A deduction rule (Φ, ϕ) is typically written as $\frac{\Phi}{\phi}$. An *axiom* is a deduction rule with an empty set of premises. We call a deduction rule *f-defining* when the outermost function symbol appearing in the source of its conclusion is f .

In this paper, for each constant c , we assume that each c -defining deduction rule is an axiom of the form $c \xrightarrow{l} p$ for some label l and closed term p .

The meaning of a TSS is defined by the following notion of least three-valued stable model. To define this notion, we need two auxiliary definitions, namely provable transition rules and contingency, which are given below.

Definition 3 (Provable Transition Rules) A deduction rule is called a transition rule when it is of the form $\frac{N}{\phi}$ with N a set of negative formulae. A TSS \mathcal{T} proves $\frac{N}{\phi}$, denoted by $\mathcal{T} \vdash \frac{N}{\phi}$, when there is a well-founded upwardly branching tree with formulae as nodes and of which

- the root is labelled by ϕ ;
- if a node is labelled by ψ and the labels of the nodes above it form the set K then:
 - ψ is a negative formula and $\psi \in N$, or
 - ψ is a positive formula and $\frac{K}{\psi}$ is a substitution instance of a deduction rule in \mathcal{T} .

Definition 4 (Contradiction, Contingency and Entailment) The formula $t \xrightarrow{l} t'$ is said to contradict $t \xrightarrow{l} \neg$, and vice versa. For two sets Φ and Ψ of formulae, Φ contradicts Ψ , denoted by $\Phi \not\models \Psi$, when there is a $\phi \in \Phi$ that contradicts a $\psi \in \Psi$. The set Φ is contingent w.r.t. Ψ , denoted by $\Phi \equiv \Psi$, when Φ does not contradict Ψ . A formula ϕ entails ψ when there is a substitution σ such that $\sigma(\phi) \equiv \psi$. A set Φ entails a set Ψ of formulae, when there exists a substitution σ such that, for each $\psi \in \Psi$, there exists a $\phi \in \Phi$ such that $\sigma(\phi) \equiv \psi$.

It immediately follows from the above definition that contradiction and contingency are symmetric relations on (sets of) formulae. We now have all the necessary ingredients to define the semantics of TSSs in terms of three-valued stable models.

Definition 5 (The Least Three-Valued Stable Model) *The pair (C, U) of sets of positive closed transition formulae is called a three-valued stable model for a TSS \mathcal{T} when*

- for each $\phi \in C$, $\mathcal{T} \vdash \frac{N}{\phi}$ for a set N of negative formulae such that $C \cup U \vDash N$,
and
- for each $\phi \in U$, $\mathcal{T} \vdash \frac{N}{\phi}$ for a set N of negative formulae such that $C \vDash N$.

C stands for Certainly and U for Unknown; the third value is determined by the formulae not in $C \cup U$. The least three-valued stable model is a three-valued stable model that is the least one with respect to the ordering on pairs of sets of formulae defined as $(C, U) \leq (C', U')$ iff $C \subseteq C'$ and $U' \subseteq U$. We say that \mathcal{T} is complete when for its least three-valued stable model it holds that $U = \emptyset$. In a complete TSS, we say that a closed substitution σ satisfies a set of formulae Φ if $\sigma(\phi) \in C$, for each positive formula $\phi \in \Phi$, and $C \vDash \sigma(\phi)$, for each negative formula $\phi \in \Phi$.

Definition 6 (Bisimulation and Bisimilarity) *Let \mathcal{T} be a TSS with signature Σ and label set \mathcal{L} . A relation $\mathcal{R} \subseteq \mathbb{C}(\Sigma) \times \mathbb{C}(\Sigma)$ is a bisimulation relation if and only if \mathcal{R} is symmetric and, for all $p_0, p_1, p'_0 \in \mathbb{C}(\Sigma)$ and $l \in \mathcal{L}$,*

$$(p_0 \mathcal{R} p_1 \wedge \mathcal{T} \vdash p_0 \xrightarrow{l} p'_0) \Rightarrow \exists p'_1 \in \mathbb{C}(\Sigma) (\mathcal{T} \vdash p_1 \xrightarrow{l} p'_1 \wedge p'_0 \mathcal{R} p'_1).$$

Two terms $p_0, p_1 \in \mathbb{C}(\Sigma)$ are called bisimilar, denoted by $p_0 \Leftrightarrow p_1$, when there exists a bisimulation relation \mathcal{R} such that $p_0 \mathcal{R} p_1$.

Bisimilarity is extended to open terms by requiring that $s, t \in \mathbb{T}(\Sigma)$ are bisimilar when $\sigma(s) \Leftrightarrow \sigma(t)$ for each closed substitution $\sigma : V \rightarrow \mathbb{C}(\Sigma)$.

3 Rule Format

We now proceed to define our rule format guaranteeing that certain constants in the language under consideration are left or right units for some binary operators. In the definition of the format proposed in the remainder of this section, we make use of a syntactic characterization of equivalence of terms up to their composition with unit elements; we call such terms *unit-context equivalent*. Intuitively, if s is unit-context equivalent to t , then s and t are bisimilar because one can be obtained from the other by applying axioms stating that some constant is a left or right unit for some binary operator. For instance, if c_1 is a left unit for a binary operator f and c_2 is a right unit for a binary operator g , then the terms $f(c_1, g(t, c_2))$ and $g(f(c_1, t), c_2)$ are both unit-context equivalent to t and also unit-context equivalent to each other.

The following definition formalizes this intuition. (While reading the technical definition, our readers may find it useful to bear in mind that $(f, c) \in L$ means that c is a left unit for a binary operator f and $(f, c) \in R$ means that c is a right unit for f .)

Definition 7 (Unit-Context Equivalent Terms) *Given sets $L, R \subseteq \Sigma \times \Sigma$ of pairs of binary function symbols and constants, $\cong^{L,R}$ is the smallest equivalence relation satisfying the following constraints, for each $s \in \mathbb{T}(\Sigma)$:*

1. $\forall_{(f,c) \in L} s \cong^{L,R} f(c, s)$, and
2. $\forall_{(g,c) \in R} s \cong^{L,R} g(s, c)$.

We say that two terms $s, t \in \mathbb{T}(\Sigma)$ are unit-context equivalent, if $s \cong^{L,R} t$.

In what follows, we abbreviate $\cong^{L,R}$ to \cong since the sets L and R are always clear from the context.

Lemma 8 *For all $s, t \in \mathbb{T}(\Sigma)$, if $s \cong t$ then $\text{vars}(s) = \text{vars}(t)$ and $\sigma(s) \cong \sigma(t)$, for each substitution σ .*

Proof. Assuming that $s \cong t$, it follows immediately from the definition of \cong that $\text{vars}(s) = \text{vars}(t)$. It is also straightforward, by an induction on the reason for $s \cong t$, to show that $\sigma(s) \cong \sigma(t)$, for each substitution σ . \square

We are now ready to define our promised rule format for unit elements.

Definition 9 (Left- and Right-Aligned Pairs) *Given a TSS, the sets L and R of pairs of binary function symbols and constants are the largest sets satisfying the following constraints.*

1. For each $(f, c) \in L$, the following conditions hold:
 - (a) For each action $a \in \mathcal{L}$, there exists at least one deduction rule of the following form:

$$\frac{\{x_0 \xrightarrow{a_i} y_i \mid i \in I\} \cup \{x_0 \xrightarrow{a_j} \mid j \in J\} \cup \{x_1 \xrightarrow{a} y_1\}}{f(x_0, x_1) \xrightarrow{a} t'}$$

where

- i. the variables y_i, y_1, x_0 and x_1 are all pairwise distinct,
- ii. for each $j \in J$, there is no c -defining axiom with a_j as label, and
- iii. there exists a collection $\{c \xrightarrow{a_i} q_i \mid i \in I\}$ of c -defining axioms such that $\sigma(t') \cong y_1$, where σ is the substitution mapping x_0 to c , each y_i to q_i , $i \in I$, and is the identity on all the other variables.

(b) Each f -defining deduction rule has the following form:

$$\frac{\Phi}{f(t_0, t_1) \xrightarrow{a} t'}$$

where $a \in \mathcal{L}$ and, for each closed substitution σ such that $\sigma(t_0) \equiv c$,

- i. either there exists some $t_1 \xrightarrow{a} t'' \in \Phi$ with $\sigma(t') \cong \sigma(t'')$, or
- ii. there exists a premise $\phi \in \Phi$ with t_0 as its source such that

- A. either ϕ is a positive formula and the collection of conclusions of c -defining deduction rules does not entail $\sigma(\phi)$, or
- B. ϕ is a negative formula and the collection of conclusions of c -defining axioms contradicts $\sigma(\phi)$.

- 2. The definition of right-aligned pairs of operators and constant symbols – that is, those such that $(f, c) \in R$ – is symmetric and is not repeated here.

For a function symbol f and a constant c , we call (f, c) left aligned (respectively, right aligned) if $(f, c) \in L$ (respectively, $(f, c) \in R$).

Condition 1a in the above definition ensures that, whenever (f, c) is in L , each transition of the form $p \xrightarrow{a} p'$, for some closed terms p and p' and action a , can be used to infer a transition $f(c, p) \xrightarrow{a} q'$ for some q' that is bisimilar to p' . This means that if (f, c) is in L then, in the context of the constant c , f does not “prune away” any of the behaviour of its first argument.

Condition 1(b)i, on the other hand, ensures that, whenever (f, c) is in L , each transition $f(c, p) \xrightarrow{a} q'$ is due to a transition $p \xrightarrow{a} p'$ for some p' that is bisimilar to q' . Thus, if (f, c) is in L then, in the context of the constant c , a term of the form $f(c, p)$ can only mimic the behaviour of p . As will become clear from the examples to follow, condition 1(b)ii ensures that the f -defining rule cannot be used to derive a transition for $f(c, p)$ and hence it is exempted from further constraints; the presence of this condition enhances the generality of our format and allows us to handle common examples of unit constants from the literature (see, e.g., Example 3).

Remark 1. Note that the requirement that $\sigma(t') \cong y_1$ in condition 1a of the above definition implies that $\text{vars}(\sigma(t')) = \{y_1\}$. Therefore x_1, y_1 and the $y_i, i \in I$, are the only variables that may possibly occur in t' .

Note that, since the sets L and R are defined as the *largest* sets of pairs satisfying the constraints from Definition 9, in order to show that (f, c) is a left-aligned pair, say, it suffices only to exhibit two sets L and R satisfying these conditions, such that (f, c) is contained in L .

The following two examples illustrate that it is in general advantageous to consider sets of left- and/or right-aligned operators instead of just a single one.

Example 1. Assume that a is the only action and consider the binary operators $f_i, i \geq 0$, with rules

$$\frac{x_1 \xrightarrow{a} y_1}{f_i(x_0, x_1) \xrightarrow{a} f_{i+1}(x_0, y_1)} .$$

Let $\mathbf{0}$ be a constant with no rules. Then each of the pairs $(f_i, \mathbf{0})$ is left aligned because the sets $L = \{(f_i, \mathbf{0}) \mid i \geq 0\}$ and $R = \emptyset$ meet the conditions from Definition 9. In particular, note that $f_{i+1}(x_0, y_1)[x_0 \mapsto \mathbf{0}] \equiv f_{i+1}(\mathbf{0}, y_1) \cong y_1$, for each $i \geq 0$. Note that, for each $i \geq 0$, the equations $f_i(\mathbf{0}, x) = x$ hold modulo bisimilarity. This fact can be checked directly by showing that the symmetric closure of the relation $\mathcal{R} = \{(f_i(\mathbf{0}, p), p) \mid p \text{ a closed term}\}$ is a bisimulation, and is also a consequence of Theorem 10 to follow, which states the correctness of the rule format we described in Definition 9.

Example 2. Consider the following TSS, which is defined for a signature with $\mathbf{0}$ and a as constants and f and g as binary function symbols.

$$\frac{}{a \xrightarrow{a} \mathbf{0}} \quad \frac{y \xrightarrow{a} y'}{f(x, y) \xrightarrow{a} g(y', x)} \quad \frac{x \xrightarrow{a} x'}{g(x, y) \xrightarrow{a} f(y, x')}$$

The TSS fits our rule format with $L = \{(f, \mathbf{0})\}$ and $R = \{(g, \mathbf{0})\}$. Note that it is essential for the above example to consider both L and R simultaneously.

Theorem 10 *Let \mathcal{T} be a complete TSS in which each rule is f -defining for some function symbol f . Assume that L and R are the sets of left- and right-aligned function symbols according to Definition 9. For each $(f, c) \in L$, it holds that $f(c, x) \leftrightarrow x$. Symmetrically, for each $(f, c) \in R$, it holds that $f(x, c) \leftrightarrow x$.*

Note that Theorem 10 trivially extends to any notion of behavioural equivalence weaker than bisimilarity.

We have already offered some intuition for the constraints in Definition 9 above. (See the paragraph following that definition.) We discuss some of the constraints in a slightly more technical fashion in Appendix A.

4 Applications and extensions

Apart from its correctness, the acid test for the usefulness of a rule format is that it be expressive enough to cover examples from the literature that afford the property they were designed to ensure. Our order of business in this section will be to offer examples of applications of the format for unit elements we introduced in Definition 9 and to show how the format can be extended to deal with operators whose semantic definition involves the use of predicates.

4.1 Applications of the basic rule format

We start by presenting examples of applications of the format for unit elements we introduced in Definition 9.

Example 3 (Nondeterministic Choice). Consider the classic nondeterministic choice operator from Milner's CCS [10] specified by the rules below, where $a \in \mathcal{L}$.

$$\frac{x \xrightarrow{a} x'}{x + y \xrightarrow{a} x'} \quad \frac{y \xrightarrow{a} y'}{x + y \xrightarrow{a} y'}$$

The sets $R = L = \{(+, \mathbf{0})\}$ meet the constraints in Definition 9. Indeed, constraint 1a and its symmetric version are trivially satisfied by the right-hand and the left-hand rule schemas, respectively. (Note that the substitution σ associated with the empty collection of axioms in condition 1(a)iii is the identity function over the set of variables.) To see that constraint 1b is also met, let σ be a closed substitution such that $\sigma(x) = \mathbf{0}$. Observe that

- each instance of the right-hand rule schema meets condition 1(b)i and
- each instance of the left-hand rule schema meets condition 1(b)iiA because the set of rules for $\mathbf{0}$ is empty and therefore does not entail $\sigma(x) = \mathbf{0} \xrightarrow{a} \sigma(x')$.

The reasoning for constraint 2 is symmetric. Therefore, Theorem 10 yields the soundness of the well known equations [8]: $\mathbf{0} + x = x = x + \mathbf{0}$.

Example 4 (Synchronous Parallel Composition). Assume, for the sake of simplicity, that a is the only action. Consider a constant RUN_a and the synchronous parallel composition from CSP [9]⁴ specified by the rules

$$\frac{}{\text{RUN}_a \xrightarrow{a} \text{RUN}_a} \quad \frac{x \xrightarrow{a} x' \quad y \xrightarrow{a} y'}{x \parallel_a y \xrightarrow{a} x' \parallel_a y'}$$

Take $L = R = \{(\parallel_a, \text{RUN}_a)\}$. These sets L and R meet the constraints in Definition 9. To see that constraint 1a and its symmetric version are satisfied by the above rule for \parallel_a , observe that the substitution σ associated with the singleton set containing the only axiom for RUN_a in condition 1(a)iii maps both the variables x and x' to RUN_a and is the identity function over the other variables. For such a σ ,

$$\sigma(x' \parallel_a y') = \text{RUN}_a \parallel_a y' \cong y'$$

To see that constraint 1b is also met, let σ be a closed substitution mapping x to RUN_a , and assume that $\text{RUN}_a \xrightarrow{a} \text{RUN}_a$ entails $\text{RUN}_a \xrightarrow{a} \sigma(x')$. It follows that $\sigma(x') = \text{RUN}_a$. Therefore,

$$\sigma(x' \parallel_a y') = \text{RUN}_a \parallel_a \sigma(y') \cong \sigma(y')$$

and constraint 1(b)i is met.

Theorem 10 thus yields the soundness of the well known equations $\text{RUN}_a \parallel_a x = x = x \parallel_a \text{RUN}_a$. These are just equation L3B from [9, page 69] and its symmetric counterpart.

Example 5 (Left Merge and Interleaving Parallel Composition). The following rules describe the operational semantics of the classic left merge and interleaving parallel composition operators [5, 10].

$$\frac{x \xrightarrow{a} x'}{x \ll y \xrightarrow{a} x' \parallel y} \quad \frac{x \xrightarrow{a} x'}{x \parallel y \xrightarrow{a} x' \parallel y} \quad \frac{y \xrightarrow{a} y'}{x \parallel y \xrightarrow{a} x \parallel y'}$$

⁴ In [9], Hoare uses the symbol \parallel to denote the synchronous parallel composition operator. Here we will use that symbol for parallel composition.

Take $L = \{(\parallel, \mathbf{0})\}$ and $R = \{(\parallel, \mathbf{0}), (\llbracket, \mathbf{0})\}$. It is easy to see that these sets L and R meet the constraints in Definition 9. Therefore, Theorem 10 yields the well known equalities

$$\mathbf{0} \parallel x = x, \quad x \parallel \mathbf{0} = x, \quad \text{and} \quad x \llbracket \mathbf{0} = x .$$

Note that the pair $(\llbracket, \mathbf{0})$ cannot be added to L while preserving constraint 1a in Definition 9. Indeed, $\mathbf{0}$ is not a left unit for the left merge operator \llbracket .

Example 6 (External Choice). The external choice operator from [9] is given by the following deduction rules

$$\frac{x \xrightarrow{a} x'}{x \square y \xrightarrow{a} x'} \quad \frac{y \xrightarrow{a} y'}{x \square y \xrightarrow{a} y'} \quad \frac{x \xrightarrow{\tau} x'}{x \square y \xrightarrow{\tau} x' \square y} \quad \frac{y \xrightarrow{\tau} y'}{x \square y \xrightarrow{\tau} x \square y'} .$$

With $L = R = \{(\square, \mathbf{0})\}$, the constraints in Definition 9 are met. The reasoning is similar to the one employed in the examples of the nondeterministic choice and synchronous parallel composition. Therefore, modulo bisimilarity, we have $\mathbf{0} \square x = x = x \square \mathbf{0}$.

Example 7 (Disrupt). Consider the following disrupt operator \blacktriangleright [3] with rules

$$\frac{x \xrightarrow{a} x'}{x \blacktriangleright y \xrightarrow{a} x' \blacktriangleright y} \quad \frac{y \xrightarrow{a} y'}{x \blacktriangleright y \xrightarrow{a} y'}$$

Note that the equation $\mathbf{0} \blacktriangleright x = x$ holds modulo bisimilarity. We now argue that its soundness is a consequence of Theorem 10. Indeed, take $L = \{(\blacktriangleright, \mathbf{0})\}$ and $R = \emptyset$. It is easy to see that these sets L and R meet the constraints in Definition 9. In particular, to see that constraint 1b is met by the first rule, observe that the set of rules for $\mathbf{0}$ is empty and therefore does not entail $\mathbf{0} \xrightarrow{a} p$ for any closed term p . A symmetric reasoning shows that the valid equation $x \blacktriangleright \mathbf{0} = x$ is also a consequence of Theorem 10.

4.2 Predicates

In the literature concerning the theory of rule formats for SOS (especially, the work devoted to congruence formats for various notions of bisimilarity), most of the time predicates are neglected at first and are only added to the considerations at a later stage. The reason is that one can encode predicates quite easily by means of transition relations. One can find a number of such encodings in the literature – see, for instance, [6, 15]. In each of these encodings, a predicate P is represented as a transition relation \xrightarrow{P} (assuming that P is a fresh label) with some fixed target. However, choosing the “right” target term to cope with the examples in the literature (and the new ones appearing in the future) within our format is extremely intricate, if not impossible. That is why we introduce an extension of our rule format that handles predicates as first-class objects, rather than coding them as transitions with dummy targets. To this end, we extend the basic notions presented in Section 2 to a setting with predicates.

Definition 11 (Predicates) Given a set \mathcal{P} of predicate symbols, Pt is a positive predicate formula and $\neg Pt$ is a negative predicate formula, for each $P \in \mathcal{P}$ and $t \in \mathbb{T}(\Sigma)$. We call t the source of both predicate formulae. In the extended setting, a (positive, negative) formula is either a (positive, negative) transition formula or (positive, negative) predicate formula. The notions of deduction rule, TSS, provable transition rules and three-valued stable models are then naturally extended by adopting the more general notion of formulae. The label of a deduction rule is either the label of the transition formula or of the predicate formula in its conclusion.

Next, we define the extension of our rule format to cater for predicates. As we did in the earlier developments, in this section we assume that, for each constant c , each c -defining deduction rule for predicates is an axiom of the form Pc .

Definition 12 (Extended Left- and Right-Aligned Pairs) Given a TSS, the sets L and R of pairs of binary function symbols and constants are the largest sets satisfying the following constraints.

1. For each $(f, c) \in L$, the following conditions hold:
 - (a) For each action $a \in \mathcal{L}$, there exists a deduction rule of the following form:

$$\frac{\{x_0 \xrightarrow{a_i} y_i \mid i \in I\} \cup \{P_k x_0 \mid k \in K\} \cup \{x_0 \xrightarrow{a_j} \text{ or } \neg P_j x_0 \mid j \in J\} \cup \{x_1 \xrightarrow{a} y_1\}}{f(x_0, x_1) \xrightarrow{a} t'}$$

where

- i. the variables y_i, y_1, x_0 and x_1 are all pairwise distinct,
 - ii. for each $j \in J$, there is no c -defining deduction rule with a_j or P_j as label (depending on whether the formula with index j is a transition or a predicate formula),
 - iii. there exists a collection $\{P_k c \mid k \in K\}$ of c -defining axioms, and
 - iv. there exists a collection $\{c \xrightarrow{a_i} q_i \mid i \in I\}$ of c -defining axioms such that $\sigma(t') \cong y_1$, where σ is the substitution mapping x_0 to c , each y_i to q_i , $i \in I$, and is the identity on all the other variables.
- (b) For each predicate $P \in \mathcal{P}$, there exists a deduction rule, of the following form:

$$\frac{\{P_i x_0 \mid i \in I\} \cup \{\neg P_j x_0 \mid j \in J\} \cup \{P x_1\}}{P f(x_0, x_1)}$$

where

- i. for each $j \in J$, there is no c -defining deduction rule with P_j as label, and
 - ii. there exists a collection $\{P_i c \mid i \in I\}$ of c -defining axioms.
- (c) Each f -defining deduction rule has one of the following forms:

$$\frac{\Phi}{f(t_0, t_1) \xrightarrow{a} t'} \quad \text{or} \quad \frac{\Phi}{P f(t_0, t_1)}$$

where $a \in \mathcal{L}$, $P \in \mathcal{P}$ and , for each closed substitution σ such that $\sigma(t_0) \equiv c$,

- i. either there exists some $t_1 \xrightarrow{a} t'' \in \Phi$ with $\sigma(t') \cong \sigma(t'')$ (if the conclusion is a transition formula), or $P t_1 \in \Phi$ (if the conclusion is a predicate formula), or
 - ii. there exists a premise $\phi \in \Phi$ with t_0 as its source such that
 - A. either ϕ is a positive formula and the collection of conclusions of c -defining deduction rules does not entail $\sigma(\phi)$, or
 - B. ϕ is a negative formula and the collection of conclusions of c -defining axioms contradicts $\sigma(\phi)$.
2. The definition of right-aligned pairs of operators and constant symbols – that is, those such that $(f, c) \in R$ – is symmetric and is not repeated here.

The definition of bisimulation is extended to a setting with predicates in the standard fashion. In particular, bisimilar terms must satisfy the same predicates.

We are now ready to state the counterpart of Theorem 10 in a setting with predicates.

Theorem 13 *Let \mathcal{T} be a complete TSS in which each rule is f -defining for some function symbol f . Assume that L and R are the sets of extended left- and right-aligned function symbols according to Definition 12. For each $(f, c) \in L$, it holds that $f(c, x) \Leftrightarrow x$. Symmetrically, for each $(f, c) \in R$, it holds that $f(x, c) \Leftrightarrow x$.*

We now provide an example of the application of the rule format offered in Definition 13. In Appendix D we give two additional examples involving the use of predicates.

Example 8 (Sequential Composition). A standard operator whose operational semantics can be given using predicates is that of sequential composition. Consider the following deduction rules, where $p \downarrow$ means that “ p can terminate successfully.” (As usual in the literature, we write the termination predicate \downarrow in postfix notation.)

$$\frac{}{1 \downarrow} \quad \frac{x \downarrow \quad y \downarrow}{x \cdot y \downarrow} \quad \frac{x \xrightarrow{a} x'}{x \cdot y \xrightarrow{a} x' \cdot y} \quad \frac{x \downarrow \quad y \xrightarrow{a} y'}{x \cdot y \xrightarrow{a} y'}$$

Take $L = R = \{(\cdot, 1)\}$. The TSS conforms to our extended rule format. The second deduction rule matches criteria 1b and 1c of Definition 12 (and the symmetric ones omitted for the right-aligned operators). The third deduction rule satisfies criterion 1(c)iiA of Definition 12 (and the omitted 2(a) and 2(c) constraints). The rightmost deduction rule satisfies constraints 1a and 1(c)i of Definition 12, as well as the omitted constraint 2(c)iiA because 1 has no transitions.

5 Conclusions

In this paper, we proposed a rule format for Structural Operational Semantics, guaranteeing constants to be left- or right-unit elements of certain operators.

The rule format encompasses advanced features such as negative premises and complex terms appearing nearly anywhere in the deduction rules. We further extended the proposed format to accommodate predicates, which are among the common ingredients in the SOS of many contemporary process description languages. The rule format is applied to a number of examples from the literature, motivating its applicability.

A straightforward extension of our rule format allows one to deal with unit elements that are complex closed terms (instead of constants). We are not aware of many practical examples in which such unit elements are present. Another algebraic property, which can be captured using the same technique, is the existence of a (left or right) zero element, i.e., a constant $\mathbf{0}$ such that $f(\mathbf{0}, x) = f(x, \mathbf{0}) = \mathbf{0}$. Mechanizing the existing rule formats for algebraic properties in a tool-set is another direction for future work.

References

1. L. Aceto, A. Birgisson, A. Ingólfssdóttir, M.R. Mousavi, and M. A. Reniers. Rule formats for determinism and idempotence. In *Proc. of FSEN'09*, Springer, 2009. To appear.
2. L. Aceto, W. J. Fokkink, and C. Verhoef. Structural operational semantics. In *Handbook of Process Algebra, Chapter 3*, pages 197–292. Elsevier, 2001.
3. J. C. M. Baeten and J. Bergstra. Mode transfer in process algebra. Tech. Rep. CSR-0001, Eindhoven University of Technology, 2000
4. J. C. M. Baeten, M. R. Mousavi, and M. A. Reniers. Timing the untimed: Terminating successfully while being conservative. In vol. 3838 of *LNCS*, pages 251–279. Springer, 2005.
5. J. A. Bergstra and J. W. Klop. Fixedpoint semantics in process algebra. Tech. Rep. IW 206/82, Center for Mathematics, Amsterdam, 1982.
6. S. Cranen, M. R. Mousavi, and M. A. Reniers. A rule format for associativity. In vol. 5201 of *LNCS*, pages 447–461, Springer, 2008.
7. R. J. van Glabbeek. The linear time - branching time spectrum I. In *Handbook of Process Algebra, Chapter 1*, pages 3–100. Elsevier, 2001.
8. M. Hennessy and R. Milner. Algebraic laws for non-determinism and concurrency. *J. ACM*, 32(1):137–161, 1985.
9. C. A. R. Hoare. *Communicating Sequential Processes*. Prentice Hall, 1985.
10. R. Milner. *Communication and Concurrency*. Prentice Hall, 1989.
11. M. R. Mousavi, M. A. Reniers, and J. F. Groote. A syntactic commutativity format for SOS. *IPL*, 93:217–223, Mar. 2005.
12. M. R. Mousavi, M. A. Reniers, and J. F. Groote. SOS formats and meta-theory: 20 years after. *TCS*, 373(3):238–272, 2007.
13. G. D. Plotkin. A structural approach to operational semantics. *JLAP*, 60-61:17-140, 2004.
14. G. D. Plotkin. A powerdomain for countable non-determinism (extended abstract). In vol. 140 of *LNCS*, pages 418–428. Springer, 1982.
15. C. Verhoef. A congruence theorem for structured operational semantics with predicates and negative premises. *Nordic Journal of Computing*, 2(2):274–302, 1995.

A Constraints of the rule format in Definition 9

Below we shall discuss some of the constraints in Definition 9 in a slightly more technical fashion. In what follows, we focus on the conditions that left-aligned pairs must meet.

First of all, note that relaxing the requirement that $x_0 \neq x_1$ in condition 1a would jeopardize Theorem 10. To see this, consider the TSS with constants $\mathbf{0}$ and a , and binary operator f with rules

$$\frac{}{a \xrightarrow{a} \mathbf{0}} \quad \frac{x \xrightarrow{a} y}{f(x, x) \xrightarrow{a} y} .$$

It is easy to check that $L = \{(f, \mathbf{0})\}$ satisfies all the constraints in condition 1 apart from $x_0 \neq x_1$. However, the term $f(\mathbf{0}, a)$ affords no transition unlike a . Therefore $\mathbf{0}$ is not a left unit for f .

In general, the requirement that variables be all different in condition 1a is needed in all the congruence rule formats for bisimilarity – see, for instance, the survey papers [2, 12] for an overview of such formats.

The role played by requirements 1(a)ii and 1(a)iii in ensuring that, modulo bisimilarity, $f(c, p)$ affords “the same transitions as p ”, for each p , is highlighted by the following two examples.

Example 9. Consider the TSS with constants $\mathbf{0}$ and a , and a binary operator f with rules:

$$\frac{}{a \xrightarrow{a} \mathbf{0}} \quad \frac{x_0 \xrightarrow{a} \quad x_1 \xrightarrow{a} y_1}{f(x_0, x_1) \xrightarrow{a} y_1} .$$

It is easy to check that $L = \{(f, a)\}$ satisfies all the constraints in condition 1 apart from 1(a)ii. However, the term $f(a, a)$ affords no transition unlike a . Therefore a is not a left unit for f .

Example 10. Consider the TSS with constants $\mathbf{0}$ and a , and a binary operator f with rules:

$$\frac{}{a \xrightarrow{a} \mathbf{0}} \quad \frac{x_0 \xrightarrow{a} y_0, \quad x_1 \xrightarrow{a} y_1}{f(x_0, x_1) \xrightarrow{a} y_1} .$$

It is easy to check that $L = \{(f, \mathbf{0})\}$ satisfies all the constraints in condition 1 apart from 1(a)iii. However, the term $f(\mathbf{0}, a)$ affords no transition unlike a . Therefore $\mathbf{0}$ is not a left unit for f .

Removing constraint 1(b)i would also jeopardize Theorem 10. To see this, consider the TSS with constant $\mathbf{0}$ and binary operator f with rules

$$\frac{x_1 \xrightarrow{a} y_1}{f(x_0, x_1) \xrightarrow{a} y_1} \quad \frac{}{f(x_0, x_1) \xrightarrow{a} \mathbf{0}} .$$

It is easy to check that $L = \{(f, \mathbf{0})\}$ satisfies all the constraints in condition 1a. However, the term $f(\mathbf{0}, \mathbf{0})$ affords an a -labelled transition unlike $\mathbf{0}$. Therefore $\mathbf{0}$ is not a left unit for f .

As we remarked previously, the role of requirement 1(b)ii is to enhance the generality of our format.

B Proof of Theorem 10

Proof. We prove that \cong is a bisimulation relation. The claim then follows since $f(c, p) \cong p$ and $g(p, c') \cong p$ for each closed term p , $(f, c) \in L$ and $(g, c') \in R$. We prove this statement by an induction on the definition of \cong . The cases that $p \cong q$ is due to reflexivity, symmetry and transitivity of \cong are trivial. So, two relevant cases remain to be proven.

1. Suppose that $p \cong q$ is due to $q \equiv f(c, p)$ for some $(f, c) \in L$.
 - (a) Assume that $p \xrightarrow{a} p' \in C$, for some $p' \in \mathbb{C}(\Sigma)$. We shall show that there exists a $p'' \in \mathbb{C}(\Sigma)$ such that $q \equiv f(c, p) \xrightarrow{a} p''$ and $p' \cong p''$.
From constraint 1a in Definition 9, we have that there exists a deduction rule of the following form

$$\frac{\{x_0 \xrightarrow{a_i} y_i \mid i \in I\} \cup \{x_0 \xrightarrow{a_j} \mid j \in J\} \cup \{x_1 \xrightarrow{a} y_1\}}{f(x_0, x_1) \xrightarrow{a} t'}$$

where

- i. the variables y_i, y_1, x_0 and x_1 are all pairwise distinct,
- ii. for each $j \in J$, there is no c -defining axiom with a_j as label, and
- iii. there exists a collection $\{c \xrightarrow{a_i} q_i \mid i \in I\}$ of c -defining axioms such that $\sigma(t') \cong y_1$, where σ is the substitution mapping x_0 to c , each y_i to q_i , $i \in I$, and is the identity on all the other variables.

Define a substitution σ' such that $\sigma'(x_0) \equiv c$, $\sigma'(x_1) \equiv p$, $\sigma'(y_1) \equiv p'$ and $\sigma'(y_i) \equiv \sigma(y_i)$ for each $i \in I$. Note that σ' satisfies all the premises in the above-mentioned rule. Therefore a proof tree for $q \equiv f(c, p) \xrightarrow{a} \sigma'(t')$ is completed. Since $\sigma(t') \cong y_1$, the last claim in Lemma 8 yields that

$$\sigma'(t') \cong \sigma'(y_1) \equiv p' .$$

Hence, $p' \cong \sigma'(t')$ and we are done.

- (b) Assume that $q \equiv f(c, p) \xrightarrow{a} q' \in C$, for some $q' \in \mathbb{C}(\Sigma)$.

By the proviso of the theorem, the transition $q \equiv f(c, p) \xrightarrow{a} q' \in C$ must be proved using an f -defining rule. Therefore, it follows from constraint 1b in Definition 9 that the transition of $q \equiv f(c, p) \xrightarrow{a} q'$ is due to a deduction rule of the following form

$$\frac{\Phi}{f(t_0, t_1) \xrightarrow{a} t'}$$

and a closed substitution σ such that $\sigma(t_0) = c$, $\sigma(t_1) = p$, $\sigma(t') = q'$ and σ satisfies Φ .

Since σ satisfies Φ and $\sigma(t_0) \equiv c$, item 1(b)ii of Definition 9 cannot apply. (Otherwise, either item 1(b)iiA holds, which means that there exists $\phi \in \Phi$ such that $\sigma(\phi) \notin C$, or item 1(b)iiB holds, which means that C contradicts $\sigma(\phi)$, or in other words, $C \vDash \sigma(\phi)$ does not hold. Any of these two implies that $\sigma(f(t_0, t_1) \xrightarrow{a} t') \in C$ is not provable using the above-given rule and the substitution σ .)

Thus according to item 1(b)i of Definition 9, there exists some $\phi \equiv t_1 \xrightarrow{a} t'' \in \Phi$ such that $q' \equiv \sigma(t') \cong \sigma(t'')$. By applying σ to ϕ , we obtain $p \equiv \sigma(t_1) \xrightarrow{a} \sigma(t'') \in C$ and we are done.

2. Suppose that $p \cong q$ is due to $q \equiv g(p, c)$ for some $(g, c) \in R$.

This case is similar to the previous case and we omit the details. \boxtimes

C Proof of Theorem 13

Proof. We prove that \cong is a bisimulation relation. The claim then follows since $f(c, p) \cong p$ and $g(p, c') \cong p$ for each closed term p , $(f, c) \in L$ and $(g, c') \in R$. We prove this statement by an induction on the definition of \cong . The cases that $p \cong q$ is due to reflexivity, symmetry and transitivity of \cong are trivial. So, two relevant cases remain to be proven.

1. Suppose that $p \cong q$ is due to $q \equiv f(c, p)$ for some $(f, c) \in L$.

- (a) Assume that $p \xrightarrow{a} p' \in C$, for some $p' \in \mathbb{C}(\Sigma)$. We shall show that there exists a $p'' \in \mathbb{C}(\Sigma)$ such that $f(c, p) \xrightarrow{a} p''$ and $p' \cong p''$.

From constraint 1a in Definition 12, we have that there exists a deduction rule of the following form

$$\frac{\{x_0 \xrightarrow{a_i} y_i \mid i \in I\} \cup \{P_k x_0 \mid k \in K\} \cup \{x_0 \xrightarrow{a_j} \text{ or } \neg P_j x_0 \mid j \in J\} \cup \{x_1 \xrightarrow{a} y_1\}}{f(x_0, x_1) \xrightarrow{a} t'}$$

where

- i. the variables y_i , y_1 , x_0 and x_1 are all pairwise distinct,
- ii. for each $j \in J$, there is no c -defining deduction rule with a_j or P_j as label (depending on whether the formula with index j is a transition or a predicate formula),
- iii. there exists a collection $\{P_k c \mid k \in K\}$ of c -defining axioms, and
- iv. there exists a collection $\{c \xrightarrow{a_i} q_i \mid i \in I\}$ of c -defining axioms such that $\sigma(t') \cong y_1$, where σ is the substitution mapping x_0 to c , each y_i to q_i , $i \in I$, and is the identity on all the other variables.

Define a substitution σ' such that $\sigma'(x_0) \equiv c$, $\sigma'(x_1) \equiv p$, $\sigma'(y_1) \equiv p'$ and $\sigma'(y_i) \equiv \sigma(y_i)$ for each $i \in I$. Note that σ' satisfies all the premises in the above-mentioned rule. Therefore a proof tree for $q \equiv f(c, p) \xrightarrow{a} \sigma'(t')$ is completed. Since $\sigma(t') \cong y_1$, the last claim in Lemma 8 yields that

$$\sigma'(t') \cong \sigma'(y_1) \equiv p' .$$

Hence, $p' \cong \sigma'(t')$ and we are done.

- (b) Assume that $q \xrightarrow{a} q' \in C$, for some $q' \in \mathbb{C}(\Sigma)$.

By the proviso of the theorem, the transition $q \equiv f(c, p) \xrightarrow{a} q' \in C$ must be proved using an f -defining rule. Therefore, it follows from constraint 1c in Definition 12 that the transition of $q \equiv f(c, p) \xrightarrow{a} q'$ is due to a deduction rule of the following form

$$\frac{\Phi}{f(t_0, t_1) \xrightarrow{a} t'}$$

and a closed substitution σ such that $\sigma(t_0) = c$, $\sigma(t_1) = p$, $\sigma(t') = q'$ and σ satisfies Φ .

Since σ satisfies Φ and $\sigma(t_0) \equiv c$, item 1(c)ii of Definition 12 cannot apply. (Otherwise, either item 1(c)iiA holds, which means that there exists $\phi \in \Phi$ such that $\sigma(\phi) \notin C$, or item 1(c)iiB holds, which means that C contradicts $\sigma(\phi)$, or in other words, $C \vDash \sigma(\phi)$ does not hold. Any of these two implies that $\sigma(f(t_0, t_1) \xrightarrow{a} t') \in C$ is not provable using the above-given rule and the substitution σ .)

Thus according to item 1(c)i of Definition 12, there exists some $\phi \equiv t_1 \xrightarrow{a} t'' \in \Phi$ such that $q' \equiv \sigma(t') \cong \sigma(t'')$. By applying σ to ϕ , we obtain $p \equiv \sigma(t_1) \xrightarrow{a} \sigma(t'') \in C$ and we are done.

- (c) Assume that $P p \in C$ for some predicate symbol $P \in \mathcal{P}$.

We proceed to show that $P q$ also holds. Since $(f, c) \in L$, condition 1b in Definition 12 yields the presence of a rule of the form

$$\frac{\{P_i x_0 \mid i \in I\} \cup \{\neg P_j x_0 \mid j \in J\} \cup \{P x_1\}}{P f(x_0, x_1)}$$

where

- i. for each $j \in J$, there is no c -defining deduction rule with P_j as label, and
- ii. there exists a collection $\{P_i c \mid i \in I\}$ of c -defining axioms.

Since $P p$ by assumption, the substitution instance of the above rule associated with any closed substitution σ mapping x_0 to c and x_1 to p proves $P f(c, p)$. Since $q \equiv f(c, p)$ we have that $P q$, which was to be shown.

- (d) Assume that $P q \in C$ for some predicate symbol $P \in \mathcal{P}$.

By the proviso of the theorem, the statement $P p \in C$ must be proved using an f -defining rule. Therefore, it follows from constraint 1c in Definition 12 that $P p$ is due to a deduction rule of the following form

$$\frac{\Phi}{P f(t_0, t_1)}$$

and a closed substitution σ such that $\sigma(t_0) \equiv c$, $\sigma(t_1) \equiv p$, and σ satisfies Φ . Since σ satisfies Φ and $\sigma(t_0) \equiv c$, item 1(c)ii of Definition 12 cannot

apply. Therefore, condition 1(c)i yields that $P t_1 \in \Phi$. Therefore $P p$ holds, as $\sigma(t_1) \equiv p$ and σ satisfies Φ .

2. Suppose that $p \cong q$ is due to $q \equiv g(p, c)$ for some $(g, c) \in R$.

This case is similar to the previous case and we omit the details. \boxtimes

D Additional examples with predicates

Example 11 (Fair Parallel Composition Operators). Consider the operators $\|_n^m$ and $\|_n$ with $m, n \geq 0$, which are inspired by those introduced in [14]. The idea is that $\|_n^m$ is a left-parallel composition for $m + 1$ steps, unless it terminates successfully before completing a sequence of $m + 1$ transitions, and then turns into $\|_n$ for some $n \geq 0$. The operator $\|_n$ behaves symmetrically.

The SOS rules for these operators are as follows, with $m, n \geq 0$.

$$\begin{array}{c}
\frac{x \downarrow \quad y \downarrow}{x \|_n^m y \downarrow} \quad \frac{x \xrightarrow{a} x'}{x \|_n^0 y \xrightarrow{a} x' \|_n y} \quad \frac{x \downarrow \quad y \xrightarrow{a} y'}{x \|_n^m y \xrightarrow{a} y'} \quad \frac{x \xrightarrow{a} x'}{x \|_n^{m+1} y \xrightarrow{a} x' \|_n^m y} \\
\frac{x \downarrow \quad y \downarrow}{x \|_n y \downarrow} \quad \frac{y \xrightarrow{a} y'}{x \|_n^0 y \xrightarrow{a} x \|_n^m y'} \quad \frac{x \xrightarrow{a} x' \quad y \downarrow}{x \|_n y \xrightarrow{a} x'} \quad \frac{y \xrightarrow{a} y'}{x \|_n^{n+1} y \xrightarrow{a} x \|_n y'}
\end{array}$$

Take $L = R = \{(\|_n^m, 1), (\|_n, 1) \mid m, n \geq 0\}$, where 1 is the constant specified in the above example. The TSS conforms to our extended rule format. By way of example, consider a pair of the form $(\|_n^m, 1) \in L$ with $m \geq 0$. The first rule for $\|_n^m$ satisfies condition 1b. The third rule for $\|_n^m$ meets requirements 1a and 1(c)i in Definition 12 for each label a . On the other hand, the second and the fourth rule meet requirement 1(c)iiA because 1 affords no transitions. The situation is entirely symmetric if we consider $(\|_n^m, 1)$ as a member of the set R .

A similar reasoning shows that the constraints in Definition 12 are met by $(\|_n, 1)$ for each $n \geq 0$.

Example 12 (Fair Parallel Composition Operators, Reprise). Consider the following variation on the above example, which uses only operators of the form $\|_n^m$ with $m \geq 0$. Again, the idea is that $\|_n^m$ is a left-parallel composition for $m + 1$ steps. However, when the “time slice” for its left argument is over, a process of the form $p \|_n^m q$ now turns into $q \|_n^n p$ for some n . Intuitively, this family of operators implements a round-robin policy with a flexible allocation of the time slice for each process.

The SOS rules for these operators are as follows, with $m, n \geq 0$.

$$\begin{array}{c}
\frac{x \downarrow \quad y \downarrow}{x \|_n^m y \downarrow} \quad \frac{x \xrightarrow{a} x'}{x \|_n^0 y \xrightarrow{a} y \|_n^n x'} \quad \frac{x \downarrow \quad y \xrightarrow{a} y'}{x \|_n^m y \xrightarrow{a} y'} \quad \frac{x \xrightarrow{a} x'}{x \|_n^{m+1} y \xrightarrow{a} x' \|_n^m y}
\end{array}$$

Take $L = R = \{(\|_n^m, 1) \mid m \geq 0\}$. Reasoning as in the previous example, it is not hard to see that the TSS conforms to our extended rule format. Note that, in order to handle this example, it is fundamental to consider the sets L and R simultaneously.